

Bosonic Part of $4d$ $N = 1$ Supersymmetric Gauge Theory with General Couplings: Local Existence

Fiki T. Akbar[#], Bobby E. Gunara^{b, #}, Triyanta^{b, #}, Freddy P. Zen^{b, #}

^b*Indonesian Center for Theoretical and Mathematical Physics (ICTMP)*

and

[#]*Theoretical Physics Laboratory*

Theoretical High Energy Physics and Instrumentation Research Group,

Faculty of Mathematics and Natural Sciences,

Institut Teknologi Bandung

Jl. Ganesha no. 10 Bandung, Indonesia, 40132

email: ft_akbar@students.itb.ac.id, bobby@fi.itb.ac.id, triyanta@fi.itb.ac.id, fpzen@fi.itb.ac.id

Abstract

In this paper, we prove the local existence of the bosonic part of $N = 1$ supersymmetric gauge theory in four dimensions with general couplings. We start with the Lagrangian of the vector and chiral multiplets with general couplings and scalar potential turned on. Then, for the sake of simplicity, we set all fermions vanish at the level of equations of motions, so we only have the bosonic parts of the theory. We apply Segal's general theory to show the local existence of solutions of equations of motions by taking Kähler potential to be bounded above by $U(n)$ symmetric Kähler potential and the first derivative of gauge couplings to be at most linear growth functions.

1 Introduction

In 1963, I. Segal has developed a method to prove the existence and uniqueness of the solutions of a semi-linear evolution equations using a semi-group [1]. Then, ten years later, he applied it to prove the existence of local and global solutions for four dimensional Yang-Mills equations in the temporal gauge condition [2]. Such a study has been extended to the case of Yang-Mills theory coupled to scalar fields in three dimensions [3], and in four dimensions [4, 5].

Our interest here is to extend the results in [4] to the case of minimal ($N = 1$) supersymmetric Yang-Mills theory coupled chiral multiplets. This theory has become a prominent subject over four decades since it might provide solutions to major problems in the Standard Model of particle physics such as the unification of gauge couplings and

the hierarchy problems.

In this paper, we prove the local existence of solutions of $N = 1$ supersymmetry gauge theory in four dimensions with general couplings. Our starting point is to consider $N = 1$ Lagrangian consists of chiral and vector with general couplings such that we have a non-linear σ -model with Kähler metric, general analytic gauge kinetic functions determined by holomorphic functions, and the scalar potential. Then, we derive field equations of motions and setting all the fermionic field to be zero at this level for the sake of simplicity. Thus, we only have an effective bosonic theory that describes the interaction between the bosonic field (ϕ, A) where ϕ is the complex scalar fields and A is the gauge fields.

By assuming that the temporal component of gauge fields vanish (analogue with the temporal gauge in Yang-Mills theory) and introducing new fields (π, E) , one can then transform the equation of motions into a semi-linear form which contains linear and non-linear terms. For our analysis, we take the fields $u = (A, E, \phi, \pi, \bar{\phi}, \bar{\pi})$ lying in $\mathcal{H} = (H_2 \times H_1)^3$ where H_p denotes a Sobolev space. We show that the linear terms are globally defined in $\mathcal{H}$ and generates a one parameter semigroup. Finally, by Segal general theory [1], the local existence of semi-linear evolution equation is established by showing that the non-linear parts satisfy local Lipschitz condition.

Since we have a generalized semi-linear evolution equation, we have to take several assumptions on the general couplings such that Segal's general theory can be used to our problem. First, we assume that Kähler potential is bounded above by $U(n)$ symmetric Kähler potential and we derive several estimates for Kähler potential and Christoffel symbol. These estimates can be used to eliminate the quantity associated with Kähler metric in our analysis. Second, we take some conditions on the gauge kinetic couplings, namely the derivative of the gauge coupling must be at most a linear growth. Finally, we assume that the scalar potential has to be at least C^3 -functions and its derivative is locally Lipschitz function.

Another problem that arises is the constraint equation which can be solved by the technique developed in [4]. This technology can be mentioned in order. We firstly decompose the E field into unique transverse (E_T) and longitudinal parts (E_L). Then by introducing a new field E_C such that $E_C = E_L$ if the constraints fulfilled, we modify the original equation of motions by replacing E_L with E_C . Using the above mentioned conditions on general couplings, we prove that the non-linear parts is locally Lipschitz function. Hence, they admit local solutions. At the end, we show that the solutions of the modified equations with the constraints satisfied are the solution of the original equation of motions.

The organization of the paper can be mentioned as follow. We shortly review a four dimensional $N = 1$ supersymmetric gauge theory in which the vector multiplets are coupled to arbitrary chiral multiplets in Section 2. Section 3 is devoted to discuss several aspects of field equation of motions including a modification of equation of motions to solve the constraint problems. In section 4, we discuss the internal scalar manifold and derive several estimates. In section 5, we prove that the non-linear part of equation of motions satisfies Lipschitz condition and finally prove the local existence.

2 General Couplings of Chiral and Vector Multiplets

In this section, we review shortly four dimensional $N = 1$ supersymmetric gauge theory in which the vector multiplets are coupled generally to arbitrary chiral multiplets. Here, we only write terms which are useful for our analysis in the paper. For an excellent review, interested reader can further consult, for example, [6, 7, 8].

The theory consists of n_v vector multiplets, (A_μ^a, λ^a) coupled to n_c chiral multiplets, (ϕ^i, χ^i) where the Latin alphabets $a, b = 1, \dots, n_v$, and $i, j = 1, \dots, n_c$ show the number of multiplets, while the Greek alphabets $\mu, \nu = 0, \dots, 3$ show the spacetime indices. In the vector multiplets we have gauge fields A_μ^a together with their fermionic partners λ^a . On the other side, the chiral multiplets contain complex scalars ϕ^i and their fermionic partners χ^i .

Furthermore, $N = 1$ supersymmetry demands the following conditions: First, the scalars ϕ^i span a Kähler manifold endowed with metric $g_{i\bar{j}} = \partial_i \partial_{\bar{j}} K$ where $K \equiv K(\phi, \bar{\phi})$ is a real function called Kähler potential. Second, there exists a set of holomorphic functions, namely (f_{ab}, X_a^i, W) which are gauge couplings, Killing vectors, and a superpotential, respectively. Finally, the existence of a real function called scalar potential which can be written as

$$V = g^{i\bar{j}} \partial_i W \partial_{\bar{j}} \bar{W} + \frac{1}{8} h^{ab} P_a P_b , \quad (2.1)$$

where P_a are real functions called Killing potentials (or momentum maps), determined by X_a^i via

$$X_a^i = \frac{i}{2} g^{i\bar{j}} \partial_{\bar{j}} P_a , \quad (2.2)$$

with h^{ab} is the inverse of $h_{ab} \equiv \text{Re} f_{ab}$. Then, one can write down the bosonic part of the $N = 1$ Lagrangian as

$$\mathcal{L} = -g_{i\bar{j}} D^\mu \phi^i D_\mu \bar{\phi}^{\bar{j}} - \frac{1}{4} h_{ab} \mathcal{F}_{\mu\nu}^a \mathcal{F}^{b\mu\nu} + \frac{1}{4} k_{ab} \mathcal{F}_{\mu\nu}^a \tilde{\mathcal{F}}^{b\mu\nu} - V , \quad (2.3)$$

where $k_{ab} \equiv \text{Im} f_{ab}$, the covariant derivative $D_\mu \phi^i \equiv \partial_\mu \phi^i + X_a^i A_\mu^a$, and the gauge field strength $\mathcal{F}_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + f_{bc}^a A_\mu^b A_\nu^c$. The dual field $\tilde{\mathcal{F}}^{a\mu\nu}$ is defined as $\tilde{\mathcal{F}}^{a\mu\nu} \equiv \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} \mathcal{F}_{\rho\sigma}^a$. It is worth to mention that Lagrangian (2.3) is invariant under the following supersymmetry transformation of the fields up to three-fermion terms

$$\begin{aligned} \delta \lambda_\bullet^a &= \frac{1}{2} \left(\mathcal{F}_{\mu\nu}^a - i \tilde{\mathcal{F}}_{\mu\nu}^a \right) \gamma^{\mu\nu} \epsilon_\bullet + 2i h^{ab} P_b \epsilon_\bullet , \\ \delta \chi^i &= i \partial_\nu \phi^i \gamma^\nu \epsilon^\bullet + 2g^{i\bar{j}} \partial_{\bar{j}} \bar{W} \epsilon_\bullet , \\ \delta A_\mu^a &= \frac{i}{2} \bar{\lambda}_\bullet^a \gamma_\mu \epsilon^\bullet + \frac{i}{2} \bar{\epsilon}_\bullet \gamma_\mu \lambda^{\bullet a} , \\ \delta \phi^i &= \bar{\chi}^i \epsilon_\bullet . \end{aligned} \quad (2.4)$$

Additionally, in the theory one can replace P_a by $P_a + \xi_a$ where ξ_a are real constants which give rise Fayet-Iliopoulos term.

In our analysis, we assume that the scalar potential to be at least a C^2 function and a satisfies the local Lipschitz condition

$$\|\partial_j V(\phi') - \partial_j V(\phi)\| \leq C(\|\phi'\|, \|\phi\|) \|\phi' - \phi\| , \quad (2.5)$$

where $C(\|\phi'\|, \|\phi\|)$ is a bounded function depend on $\|\phi\|$. The condition above implies that the holomorphic superpotential has to be at least a C^3 function.

3 Field Equations of Motions

This section is devoted to discuss several aspect of field equation of motions. In particular, we take all fermions to be trivial at this level. Thus, the gauge fields and the scalars are the main ingredients of our analysis in this paper. To simplify the analysis we take a condition $A_0 = 0$ which is not a gauge condition since in general Lagrangian (2.3) is no longer gauge invariant ¹.

First of all, the gauge field equation of motions is given by

$$D_\mu \left(h_{ab} \mathcal{F}^{b\mu\nu} - k_{ab} \tilde{\mathcal{F}}^{b\mu\nu} \right) = g_{i\bar{j}} \left(X_a^i D^\nu \bar{\phi}^{\bar{j}} + \bar{X}_a^{\bar{j}} D^\nu \phi^i \right), \quad (3.1)$$

with

$$D_\mu \left(h_{ab} \mathcal{F}^{b\mu\nu} \right) = \partial_\mu \left(h_{ab} \mathcal{F}^{b\mu\nu} \right) + f_{ac}^d A_\mu^c \left(h_{db} \mathcal{F}^{b\mu\nu} \right). \quad (3.2)$$

The field strength tensor $\mathcal{F}_{\mu\nu}^a$ satisfies Bianchi identity,

$$D_\mu \tilde{\mathcal{F}}^{b\mu\nu} = 0. \quad (3.3)$$

By defining

$$E^{as} = -\mathcal{F}^{a0s} = -\partial^0 A^{ar}, \quad (3.4)$$

and using (3.3), we can rewrite (3.1) as

$$\begin{aligned} \frac{\partial E^{ar}}{\partial t} = & \partial_s \partial^s A^{ar} - \partial^r \partial_s A^{as} + f_{bc}^a \partial_s (A^{as} A^{ar}) + h^{ab} \mathcal{F}^{csr} \partial_s h_{bc} + h^{ab} h_{de} f_{bc}^d A_s^c \mathcal{F}^{esr} \\ & - h^{ab} E^{cr} \partial_0 h_{bc} - h^{ab} \tilde{\mathcal{F}}^{c0r} \partial_0 k_{bc} - h^{ab} \tilde{\mathcal{F}}^{csr} \partial_s k_{bc} - h^{ab} (k_{de} f_{bc}^d - k_{bd} f_{ce}^d) A_s^c \mathcal{F}^{esr} \\ & - h^{ab} g_{i\bar{j}} \left(X_a^i D^r \bar{\phi}^{\bar{j}} + \bar{X}_a^{\bar{j}} D^r \phi^i \right), \end{aligned} \quad (3.5)$$

together with the constraint equations

$$\mathcal{C}^a(t) = -\partial_s E^{as} + 4\pi \rho^a, \quad (3.6)$$

where

$$\begin{aligned} 4\pi \rho^a = & h^{ab} g_{i\bar{j}} \left(X_b^i D^0 \bar{\phi}^{\bar{j}} + \bar{X}_b^{\bar{j}} D^0 \phi^i \right) + h^{ab} \tilde{\mathcal{F}}^{cs0} \partial_s k_{bc} - h^{ab} E^{cs} \partial_s h_{bc} \\ & h^{ab} (k_{de} f_{bc}^d - k_{bd} f_{ce}^d) A_s^c \tilde{\mathcal{F}}^{es0} - h^{ab} h_{de} f_{bc}^d A_s^c E^{es}, \end{aligned} \quad (3.7)$$

with the initial value of (3.6) is $\mathcal{C}^a(0) = 0$. Next, we consider the scalar field equation of motions which have been modified into

$$\begin{aligned} \frac{\partial \pi^i}{\partial t} = & \partial_s \partial^s \phi^i + \Gamma_{kl}^i \left(\partial_r \phi^k \partial^r \phi^l - \pi^k \pi^l \right) + X_a^i \partial_r A^{ar} + A^{ar} \partial_r \phi^k \nabla_k X_a^i \\ & + g^{i\bar{j}} g_{k\bar{l}} A^{ar} D_r \phi^k \nabla_{\bar{j}} \bar{X}_a^{\bar{l}} - g^{i\bar{j}} \left(\frac{1}{4} h_{ab\bar{j}} \mathcal{F}_{\mu\nu}^a \mathcal{F}^{b\mu\nu} - \frac{1}{4} k_{ab\bar{j}} \mathcal{F}_{\mu\nu}^a \tilde{\mathcal{F}}^{b\mu\nu} + \partial_{\bar{j}} V_S \right), \end{aligned} \quad (3.8)$$

where

$$\pi^i = -D^0 \phi^i, \quad (3.9)$$

$$\nabla_k X_a^i = \partial_k X_a^i + \Gamma_{kl}^i X_a^l, \quad (3.10)$$

¹For renormalizable Yang-Mills-Higgs theory with constant f_{ab} and $g_{i\bar{j}}$ the condition $A_0 = 0$ is called temporal gauge, see for example, [9].

together with its complex conjugate.

Now, we rewrite the equations of motion (3.1) and (3.8) into the following form

$$\frac{du}{dt} = \mathcal{A}u + J(u) , \quad (3.11)$$

where

$$u = \begin{bmatrix} A^{as} \\ E^{as} \\ \phi^i \\ \pi^i \\ \bar{\phi}^{\bar{j}} \\ \bar{\pi}^{\bar{j}} \end{bmatrix} , \quad \mathcal{A}u = \begin{bmatrix} E^{as} \\ \partial_r \partial^r A^{as} - \partial_r \partial^s A^{ar} \\ \pi^i \\ \partial_r \partial^r \phi^i \\ \bar{\pi}^{\bar{j}} \\ \partial_r \partial^r \bar{\phi}^{\bar{j}} \end{bmatrix} ,$$

and

$$J(u) = \begin{bmatrix} 0 \\ f_{bc}^a \partial_r (A^{bs} A^{cr}) + h^{ab} \left(\mathcal{F}^{esr} \partial_r h_{bc} - \tilde{\mathcal{F}}^{esr} \partial_r k_{bc} + h_{de} f_{bc}^d A_r^c \mathcal{F}^{esr} \right) + \mathcal{D}_1 \\ 0 \\ \Gamma_{kl}^i (\partial_s \phi^k \partial^s \phi^l - \pi^k \pi^l) + X_a^i \partial_s A^{as} + A^{as} \nabla_k X_a^i \partial_s \phi^k + g^{i\bar{j}} g_{l\bar{k}} A_s^a \nabla_{\bar{j}} \bar{X}_a^{\bar{k}} D^s \phi^l + \mathcal{D}_2 \\ 0 \\ \Gamma_{\bar{k}\bar{l}}^{\bar{j}} \partial_s \bar{\phi}^{\bar{k}} \partial^s \bar{\phi}^{\bar{l}} + \bar{X}_a^{\bar{j}} \partial_s A^{as} + A^{as} \nabla_{\bar{k}} \bar{X}_a^{\bar{j}} \partial_s \bar{\phi}^{\bar{k}} - g^{i\bar{j}} g_{kl} A_s^a \nabla_i X_a^k \left\{ \partial^s \bar{\phi}^{\bar{l}} + A^{bs} \bar{X}_b^{\bar{l}} \right\} + \bar{\mathcal{D}}_2 \end{bmatrix} . \quad (3.12)$$

where,

$$\begin{aligned} \mathcal{D}_1 &= -h^{ab} \left(h_{bci} \pi^i + h_{bc\bar{j}} \bar{\pi}^{\bar{j}} \right) E^{cs} - h^{ab} \left(k_{bci} \pi^i + k_{bc\bar{j}} \bar{\pi}^{\bar{j}} \right) \tilde{\mathcal{F}}^{c0s} \\ &\quad - h^{ab} \left(k_{de} f_{bc}^d - k_{bd} f_{ce}^d \right) A_r^c \tilde{\mathcal{F}}^{esr} - h^{ab} g_{i\bar{j}} \left(X_b^i D^s \bar{\phi}^{\bar{j}} + X_b^{\bar{j}} D^s \phi^i \right) , \end{aligned} \quad (3.13)$$

$$\mathcal{D}_2 = g^{i\bar{j}} \frac{h_{ab\bar{j}}}{4} \left(-2E_s^a E^{bs} - \mathcal{F}_{rs}^a \mathcal{F}^{brs} \right) + g^{i\bar{j}} \frac{k_{ab\bar{j}}}{2} \epsilon^{srl} E_s^a \mathcal{F}_{rl}^b - g^{i\bar{j}} \partial_{\bar{j}} V , \quad (3.14)$$

$$\bar{\mathcal{D}}_2 = g^{i\bar{j}} \frac{h_{abi}}{4} \left(-2E_s^a E^{bs} - \mathcal{F}_{rs}^a \mathcal{F}^{brs} \right) + g^{i\bar{j}} \frac{k_{abi}}{2} \epsilon^{srl} E_s^a \mathcal{F}_{rl}^b - g^{i\bar{j}} \partial_i V , \quad (3.15)$$

together with the constraint equation (3.6).

To solve the constraint problem, we use the method in [4] which can be structured as follows. First, we modify (3.1) by decomposing E field into unique transverse parts (divergence free) E_T and longitudinal parts (curl free) E_L

$$E = E_T + E_L , \quad (3.16)$$

with

$$\partial_s E_T^s = 0, \quad \epsilon_{qrs} \partial^r E_L^s = 0 . \quad (3.17)$$

Then, replacing E_L with the new fields, E_C which equals to E_L when the constraint is satisfied (see lemma 3),

$$E_L^s \rightarrow E_C^s = \partial_s \left(-\frac{1}{4\pi R} * \rho \right) , \quad (3.18)$$

where $-\frac{1}{4\pi R} * \rho$ represents convolution of ρ with the fundamental solution of Poisson equation,

$$-\frac{1}{4\pi R} * \rho = -\frac{1}{4\pi} \int_{\mathbb{R}^3} dx' \left(\frac{\rho(x')}{|x - x'|} \right) . \quad (3.19)$$

Let $\mathcal{H} = (H_2 \times H_1)^3$, where H_p represent a Sobolev space of square integrable functions over $\mathbb{R}^3$ with their derivative up to order p are also square integrable and let $\|\cdot\|_{H_p}$ represents a Sobolev norm defined as

$$\|u\|_{H_p} = \left[\sum_{|\alpha|=0}^p \|D^\alpha u\|_{L_2}^2 \right]^{\frac{1}{2}}. \quad (3.20)$$

where $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_m)$ is a multi-index of non-negative integers and $|\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_m$. The H_p space is also a Hilbert space. In our analysis, we take the fields $u = (A, E, \phi, \pi, \bar{\phi}, \bar{\pi})$ to lie in $\mathcal{H}$.

Now we prove the following lemmas which are an extension of lemmas in [4] to incorporate the gauge kinetic functions and show that for fields u lying in $\mathcal{H}$, the condition $E_L = E_C$ implies that E satisfies the constraint equation.

Lemma 1. *Let ρ is defined as in 3.7. If the gauge kinetic function is also lies at least in H_1 , then $E_C^s = \partial^s \left(-\frac{1}{4\pi r} * \rho \right) = \frac{1}{4\pi} \left(\frac{\hat{r}}{r^2} * \rho \right)^s$.*

Proof. Since $\frac{1}{r}$ is a weak $L_{3,w}(\mathbb{R}^3)$ function, then using generalize Young inequality for convolution product [10], we have

$$\left\| \frac{1}{r} * \rho \right\|_{L_p} \leq C \|\rho\|_{L_q} \left\| \frac{1}{r} \right\|_{3,w} \leq C' \|\rho\|_{L_q}, \quad (3.21)$$

for $\frac{1}{q} = \frac{1}{p} + \frac{2}{3}$. It means $\frac{1}{r} * \rho \in L_p(\mathbb{R}^3)$ provided that $\rho \in L_q(\mathbb{R}^3)$. Since all field $(A, E, \phi, \pi, \bar{\phi}, \bar{\pi})$ at least lie in H_1 and using the assumption that the gauge kinetic function is also lie at least in H_1 , then by Sobolev embedding theorem, they all lie in L_s for $2 \leq s \leq 6$. Then, $\rho \in L_q$ for $1 \leq q \leq 3$, thus $\frac{1}{r} * \rho \in L_p(\mathbb{R}^3)$ for $p > 3$.

Let V be an arbitrary C^∞ rapidly decreasing vector field in $\mathbb{R}^3$, we have

$$\begin{aligned} \langle V_s, E_C^s \rangle &= \frac{1}{4\pi} \left\langle V_s, -\partial^s \left(\frac{1}{r} * \rho \right) \right\rangle \\ &= \frac{1}{4\pi} \left\langle \partial^s V_s, \left(\frac{1}{r} * \rho \right) \right\rangle \\ &= \frac{1}{4\pi} \left\langle \frac{1}{r} * \partial^s V_s, \rho \right\rangle. \end{aligned} \quad (3.22)$$

Since $\left| \frac{\hat{r}}{r} \right|$ is a weak $L_{3/2,w}(\mathbb{R}^3)$ function and $\rho \in L_q(\mathbb{R}^3)$ for $1 \leq q \leq 3$, then

$$\left| \frac{\hat{r}}{r} \right| * |\rho| \in L_r, \quad (3.23)$$

for all $\frac{3}{2} < r < 3$. Hence, we have,

$$\begin{aligned} \frac{1}{4\pi} \left\langle \frac{1}{r} * \partial^s V_s, \rho \right\rangle &= \frac{1}{4\pi} \int V_s \left[\int \frac{x^s - x'^s}{|x - x'|^3} \rho(x') dx' \right] dx \\ &= \frac{1}{4\pi} \left\langle V_s, \left(\frac{\hat{r}}{r^2} * \rho \right)^s \right\rangle. \end{aligned} \quad (3.24)$$

And finally we have $E_C^s = \partial^s \left(-\frac{1}{4\pi r} * \rho \right) = \frac{1}{4\pi} \left(\frac{\hat{r}}{r^2} * \rho \right)^s$ which complete the proof. $\square$

Lemma 2. $E_C \in L_p$ for all $\frac{3}{2} < p < \infty$. In particular $E_C \in L_2$.

Proof. From the proof of the previous lemma, we already have $\rho \in L_q(\mathbb{R}^3)$ for $1 \leq q \leq 3$. Then using the representation of E_C in previous lemma,

$$\|E_C\|_{L_p} = \left\| \frac{\hat{r}}{r^2} * \rho \right\|_{L_p} \leq C \|\rho\|_{L_q} \left\| \frac{\hat{r}}{r^2} \right\|_{3/2, w} \leq C' \|\rho\|_{L_q}, \quad (3.25)$$

for $1 < p, q < \infty$ and $\frac{1}{q} = \frac{1}{p} + \frac{1}{3}$, then $E_C \in L_s$ for all $\frac{3}{2} < s < \infty$.

In particular, $E_C \in L_2$ and the following inequality holds,

$$\|E_C\|_{L_2} \leq C \|\rho\|_{L_{6/5}}. \quad (3.26)$$

□

Lemma 3. The condition $E_C = E_L$ is equivalent to $\partial_s E^s = 4\pi\rho$.

Proof. Let V be an arbitrary C^∞ rapidly decreasing function in $\mathbb{R}^3$, we have

$$\begin{aligned} \langle V, \partial_s E_C^s \rangle &= \frac{1}{4\pi} \int V(x) \partial_s \left[\int \frac{x^s - x'^s}{|x - x'|^3} \rho(x') dx' \right] dx \\ &= -\frac{1}{4\pi} \int \rho(x') \left[\int \frac{x^s - x'^s}{|x - x'|^3} \partial_s V dx \right] dx' \\ &= \frac{1}{4\pi} \int \rho(x') \left[\int V(x) \partial_s \left(\frac{x^s - x'^s}{|x - x'|^3} \right) dx \right] dx' \\ &= \int \rho(x') V(x') dx' \\ &= \langle V, \rho \rangle, \end{aligned} \quad (3.27)$$

where we used integrating by parts and exchange the order of integration. Thus, E_C satisfies $\partial_s E_C^s = 4\pi\rho$ as distribution.

Since $E_C \in L^2$, we can define it's Fourier transform and decomposing as

$$\hat{E}_C = \hat{E}_C^T + \hat{E}_C^L, \quad (3.28)$$

with

$$\left(\hat{E}_C^T \right)^s = \left(\delta^{sr} - \frac{k^s k^r}{|k|^2} \right) \left(\hat{E}_C \right)_r \quad (3.29)$$

$$\left(\hat{E}_C^L \right)^s = \frac{k^s k^r}{|k|^2} \left(\hat{E}_C \right)_r. \quad (3.30)$$

Furthermore, E_C is a gradient, then E_C has vanishing curl, then $\hat{k} \times \hat{E}_C = 0$ which implies that E_C has only a longitudinal component, thus

$$\left(E_C^L \right)^s = \frac{k^s k^r}{|k|^2} \left(\hat{E}_C \right)_r = \left(\hat{E}_C \right)^s, \quad (3.31)$$

and because E_C satisfies $\partial_s E_C^s = 4\pi\rho$, taking a Fourier transform, we have

$$\hat{E}_C^s = -i \frac{k^s}{|k|^2} (4\pi\hat{\rho}). \quad (3.32)$$

Now suppose that the field u satisfies the constraint equation,

$$\partial_s E^s = \partial_s E_L^s = 4\pi\rho, \quad (3.33)$$

where we used the decomposition in (3.16). Taking a Fourier transform of the constraint equation, we get

$$\begin{aligned} \hat{E}_L^s &= -i \frac{k^s}{|k|^2} (4\pi\hat{\rho}) \\ &= \hat{E}_C^s. \end{aligned} \quad (3.34)$$

Thus any solution of the constraint has $E_L = E_C$. Conversely, if $E_L = E_C$, the constraint is satisfied. $\square$

With the modification, we can rewrite the equation of motions as follows

$$\frac{du}{dt} = \mathcal{A}u + J(u), \quad (3.35)$$

where

$$u = \begin{bmatrix} A^{as} \\ E^{as} \\ \phi^i \\ \pi^i \\ \bar{\phi}^{\bar{j}} \\ \bar{\pi}^{\bar{j}} \end{bmatrix}, \quad \mathcal{A}u = \begin{bmatrix} E_T^{as} \\ \partial_r \partial^r A^{as} - \partial_r \partial^s A^{ar} \\ \pi^i \\ \partial_r \partial^r \phi^i \\ \bar{\pi}^{\bar{j}} \\ \partial_r \partial^r \bar{\phi}^{\bar{j}} \end{bmatrix},$$

and

$$J(u) = \begin{bmatrix} \partial^s \left\{ -\frac{1}{4\pi r} * \rho \right\} \\ f_{bc}^a \partial_r (A^{bs} A^{cr}) + h^{ab} \left(\mathcal{F}^{esr} \partial_r h_{bc} - \tilde{\mathcal{F}}^{esr} \partial_r k_{bc} + h_{de} f_{bc}^d A_r^c \mathcal{F}^{esr} \right) + \mathcal{D}_1 \\ 0 \\ \Gamma_{kl}^i (\partial_s \phi^k \partial^s \phi^l - \pi^k \pi^l) + X_a^i \partial_s A^{as} + A^{as} \nabla_k X_a^i \partial_s \phi^k + g^{i\bar{j}} g_{l\bar{k}} A_s^a \nabla_{\bar{j}} \bar{X}_a^{\bar{k}} D^s \phi^l + \mathcal{D}_2 \\ 0 \\ \Gamma_{\bar{k}\bar{l}}^{\bar{j}} \partial_s \bar{\phi}^{\bar{k}} \partial^s \bar{\phi}^{\bar{l}} + \bar{X}_a^{\bar{j}} \partial_s A^{as} + A^{as} \nabla_{\bar{k}} \bar{X}_a^{\bar{j}} \partial_s \bar{\phi}^{\bar{k}} - g^{i\bar{j}} g_{k\bar{l}} A_s^a \nabla_i X_a^k \left\{ \partial^s \bar{\phi}^{\bar{l}} + A^{bs} \bar{X}_b^{\bar{l}} \right\} + \bar{\mathcal{D}}_2 \end{bmatrix}. \quad (3.36)$$

4 Scalar Internal Manifold

This section is assigned for the discussion of the internal scalar manifold. In particular, we consider the case of the Kähler potential to be bounded to a function and derive an estimates for Kähler potential and Christoffel symbol. Our estimates derived in this section is important in our analysis for proving the local existence of (3.35).

As mention in section 2, in four dimensions, the $N = 1$ supersymmetry theory demands that the scalar field $(\phi, \bar{\phi})$ span a Kähler manifold with Kähler potential $K \equiv K(\phi, \bar{\phi})$. In this paper, we consider the case where the Kähler potential bounded above by $U(n_c)$ symmetric Kähler potential and satisfies several conditions,

$$K \leq \Phi(|\phi|), \quad (4.1)$$

$$|\Gamma| \leq |\tilde{\Gamma}|, \quad (4.2)$$

where $|\phi| = (\delta_{i\bar{j}}\phi^i\bar{\phi}^{\bar{j}})^{\frac{1}{2}}$ and $\tilde{\Gamma}$ is the Christoffel symbol of $\tilde{g}$.

We prove a lemma about estimates of Kähler potential and Christoffel symbol,

Lemma 4. *Let $\mathcal{M}$ be a Kähler manifold with Kahler potential $K = K(\phi, \bar{\phi})$. If $\mathcal{M}$ satisfies (4.1), (4.2) and*

$$\left| \frac{F'}{2|\phi|} \right| \leq \epsilon, \quad (4.3)$$

where $F(|\phi|) = \frac{1}{4|\phi|^2} \left(\Phi'' - \frac{\Phi'}{|\phi|} \right)$ with $\Phi' = \partial\Phi/\partial|\phi|$ and ϵ is a non negative constant, then we have the following estimates

$$|K| \leq \frac{\epsilon}{6} |\phi|^6 + \frac{C_1}{2} |\phi|^4 + C_2 |\phi|^2 + C_3, \quad (4.4)$$

$$|\Gamma| \leq 2\epsilon |\phi|^3 + C_1 |\phi|. \quad (4.5)$$

Proof. Let $\tilde{\mathcal{M}}$ be a Kähler manifold generated by Φ . We can write the metric $\tilde{g}_{i\bar{j}} = \partial_i \partial_{\bar{j}} \Phi$ as

$$\tilde{g}_{i\bar{j}} = \frac{\Phi'}{2|\phi|} \delta_{i\bar{j}} + \frac{1}{4|\phi|^2} \left(\Phi'' - \frac{\Phi'}{|\phi|} \right) \delta_{k\bar{j}} \delta_{i\bar{k}} \phi^k \bar{\phi}^{\bar{k}}, \quad (4.6)$$

where $\Phi' = \partial\Phi/\partial|\phi|$. The inverse of the metric can written as,

$$\tilde{g}^{i\bar{j}} = \frac{2|\phi|}{\Phi'} \delta^{i\bar{j}} - \frac{2}{\Phi'|\phi|} \left(\frac{\Phi'' - \frac{\Phi'}{|\phi|}}{\Phi'' + \frac{\Phi'}{|\phi|}} \right) \phi^i \bar{\phi}^{\bar{j}}. \quad (4.7)$$

The norm of the Christoffell symbol is

$$\begin{aligned} |\tilde{\Gamma}| &= \left(\tilde{g}^{j\bar{j}} \tilde{g}^{k\bar{k}} \tilde{g}_{i\bar{i}} \tilde{\Gamma}_{jk}^i \tilde{\Gamma}_{\bar{j}\bar{k}}^{\bar{i}} \right)^{\frac{1}{2}} \\ &= \left(\tilde{g}^{j\bar{j}} \tilde{g}^{k\bar{k}} \tilde{g}_{i\bar{i}} \tilde{g}^{i\bar{l}} \partial_j \tilde{g}_{k\bar{l}} \tilde{g}^{\bar{l}\bar{i}} \partial_{\bar{j}} \tilde{g}_{l\bar{k}} \right)^{\frac{1}{2}} \\ &= \left(\tilde{g}^{l\bar{l}} \tilde{g}^{j\bar{j}} \tilde{g}^{k\bar{k}} \partial_j \tilde{g}_{k\bar{l}} \partial_{\bar{j}} \tilde{g}_{l\bar{k}} \right)^{\frac{1}{2}} \\ &= |\partial \tilde{g}|, \end{aligned} \quad (4.8)$$

and the first derivative of the metric is

$$\partial_j \tilde{g}_{k\bar{l}} = F (\delta_{k\bar{l}} \delta_{j\bar{i}} + \delta_{k\bar{i}} \delta_{j\bar{l}}) \bar{\phi}^{\bar{i}} + \frac{F'}{2|\phi|} \delta_{k\bar{i}} \delta_{i\bar{l}} \delta_{j\bar{m}} \phi^i \bar{\phi}^{\bar{i}} \bar{\phi}^{\bar{m}}, \quad (4.9)$$

where

$$\begin{aligned} F(|\phi|) &= \frac{1}{4|\phi|^2} \left(\Phi'' - \frac{\Phi'}{|\phi|} \right), \\ \frac{F'}{2|\phi|} &= \frac{1}{8|\phi|^3} \left(\Phi''' - \frac{3\Phi''}{|\phi|} + \frac{3\Phi'}{|\phi|^2} \right). \end{aligned} \quad (4.10)$$

If condition (4.3) is satisfied, then using the following inequality for integral,

$$\left| \int f(x) dx \right| \leq \int |f(x)| dx, \quad (4.11)$$

we have the following estimates,

$$\begin{aligned} |F| &\leq \epsilon |\phi|^2 + C_1, \\ |\Phi| &\leq \frac{\epsilon}{6} |\phi|^6 + \frac{C_1}{2} |\phi|^4 + C_2 |\phi|^2 + C_3, \end{aligned} \quad (4.12)$$

where $C_1 = |F(0)|$, $C_2 = \left| \frac{\Phi'}{2|\phi|}(0) \right|$ and $C_3 = |\Phi(0)|$. Then we have the norm of the Christoffel symbol satisfies

$$|\tilde{\Gamma}| \leq 2\epsilon |\phi|^3 + C_1 |\phi|. \quad (4.13)$$

Hence, by our assumption in 4.1 and 4.2, we have

$$|\Phi| \leq \frac{\epsilon}{6} |\phi|^6 + \frac{C_1}{2} |\phi|^4 + C_2 |\phi|^2 + C_3, \quad (4.14)$$

$$|\Gamma| \leq 2\epsilon |\phi|^3 + C_1 |\phi|. \quad (4.15)$$

This complete the proof. $\square$

Our assumption in (4.3) is satisfied for several examples of Kähler manifold, for examples are $\mathbb{C}^n$ and $\mathbb{C}P^n$ which are widely used in the theory. For $\mathbb{C}^n$, the Kähler potential is given by $|\phi|^2$, then clearly F is vanish, hence $\frac{F'}{2|\phi|}$ is bounded by 0. In case of $\mathbb{C}P^n$, the Kähler potential (using standard Fubini-Study metric) is given by

$$\Phi_{\mathbb{C}P^n}(|\phi|) = \ln(1 + |\phi|^2). \quad (4.16)$$

Then we have,

$$F = -\frac{1}{(1 + |\phi|^2)^2}, \quad (4.17)$$

and

$$\left| \frac{F'}{2|\phi|} \right| = \frac{2}{(1 + |\phi|^2)^3}, \quad (4.18)$$

which is bounded above by 2.

5 Local Existence

In this section, we will prove the local existence of the evolution equation (3.35) using Segal's theorem. Furthermore, we shall show that solutions of (3.35) are the solutions of the original equations, namely (3.1) and (3.8).

In section 3, we have derived the equation of motions for u and have taken the field u lie in $\mathcal{H} = (H_2 \times H_1)^3$. Let us consider the linear part of the evolution equation (3.35),

$$\frac{du}{dt} = \mathcal{A}u. \quad (5.1)$$

By decomposing the A and E fields into transverse and longitudinal components, we can write (5.1) as follow,

$$\frac{d}{dt} \begin{bmatrix} A_T^{as} \\ E_T^{as} \end{bmatrix} = \begin{bmatrix} 0 & I \\ \Delta & 0 \end{bmatrix} \begin{bmatrix} A_T^{as} \\ E_T^{as} \end{bmatrix}, \quad \frac{d}{dt} \begin{bmatrix} A_L^{as} \\ E_L^{as} \end{bmatrix} = 0, \quad (5.2)$$

$$\frac{d}{dt} \begin{bmatrix} \phi^i \\ \pi^i \end{bmatrix} = \begin{bmatrix} 0 & I \\ \Delta & 0 \end{bmatrix} \begin{bmatrix} \phi^i \\ \pi^i \end{bmatrix}, \quad \frac{d}{dt} \begin{bmatrix} \bar{\phi}^{\bar{j}} \\ \bar{\pi}^{\bar{j}} \end{bmatrix} = \begin{bmatrix} 0 & I \\ \Delta & 0 \end{bmatrix} \begin{bmatrix} \bar{\phi}^{\bar{j}} \\ \bar{\pi}^{\bar{j}} \end{bmatrix}. \quad (5.3)$$

Each pair of fields (A_T, E_T) , (ϕ, π) , and $(\bar{\phi}, \bar{\pi})$ satisfies the linear wave equation and (A_L, E_L) is a constant of the linearized equation. Thus, the linear operator $\mathcal{A}$ generates a one-parameter semigroup on $\mathcal{H}$ and for any initial value $u_0 = u(t_0) \in \mathcal{H}$, the linearized equation admits a classical solution which can be written as ²,

$$u(t) = e^{\mathcal{A}(t-t_0)} u_0 . \quad (5.4)$$

Then, it follows that the solution of linearized equation is globally defined on $\mathcal{H}$.

Following the result above, by writing the evolution equation (3.35) as an integral equation,

$$u(t) = e^{\mathcal{A}(t-t_0)} u_0 + \int_{t_0}^t ds e^{\mathcal{A}(s-t_0)} J(u(s)) , \quad (5.5)$$

the local existence of solution of the equation is established by showing that the nonlinear operator J satisfies Lipschitz condition,

$$\|J(u') - J(u)\| \leq C(\|u'\|, \|u\|) \|u' - u\| , \quad (5.6)$$

for all $u', u \in \mathcal{H}$. The norm $\|\cdot\|$ is designed for $\mathcal{H}$ norm and $C(\cdot, \cdot)$ is some monotonically increasing, finite function of the norm indicated. Then, for any initial data $u_0 = u(t_0) \in D_{\mathcal{A}}$, where $D_{\mathcal{A}}$ is a domain of linear operator $\mathcal{A}$, the evolution equation (3.35) admits a unique classical solution on some interval (T_1, T_2) containing t_0 either $(T_1, T_2) = (-\infty, \infty)$ or $\|u(t)\| \rightarrow \infty$ as $t \rightarrow T_1$ or T_2 .

Let us write the components of the non-linear operator J as $J = (J_1, J_2, J_3, J_4, J_5, J_6)$. The proof that J satisfies a Lipschitz condition is facilitated by a Schauder ring property for Sobolev space over $\mathbb{R}^3$ and the Sobolev inequality over $\mathbb{R}^3$,

$$\|\partial^r u\|_{L_p} \leq C \|\partial^s u\|_{L_m}^\theta \|u\|_{L_q}^{1-\theta} , \quad (5.7)$$

for real numbers q, m with $1 \leq q, m \leq \infty$ and r, s are integers where $0 \leq r < s$ which satisfy

$$\frac{1}{p} = \frac{r}{3} + \theta \left(\frac{1}{m} - \frac{s}{3} \right) + (1 - \theta) \frac{1}{q} , \quad (5.8)$$

with $r/s \leq \theta \leq 1$ and p is non negative, and a constant C depends only on m, j, q, r and θ . For further discussion on Sobolev inequality, see [12].

Now, since the theory has scalar fields dependent gauge couplings, we have to make an assumption for the gauge kinetic function in order to prove that E_C is a mapping from $\mathcal{H}$ to H_2 and locally Lipschitz continuous.

Lemma 5. *Let the fields $(A, E, \phi, \pi, \bar{\phi}, \bar{\pi}) \in \mathcal{H}$. If the first derivative of the real part of the gauge kinetic function is at most a linear growth,*

$$\|\partial_i h_{ab}\|_{H_1} = \|\partial_{\bar{j}} h_{ab}\|_{H_1} \leq C \|\phi\|_{H_2} , \quad (5.9)$$

then E_C lies in H_2 . Furthermore, if $\partial_s h_{ab}$ is also a locally Lipschitz function, then E_C is a locally Lipschitz function.

²for details review, see [1], [11].

Proof. Define a norm for a gauge indexed field as

$$|A|^2 = h_{ab} A^{as} A_s^b . \quad (5.10)$$

By definition of Sobolev norm, we have

$$\|E_C\|_{H_2} = (\|E_C\|_{L_2}^2 + \|D^2 E_C\|_{L_2}^2)^{1/2} . \quad (5.11)$$

Using (3.26), we have

$$\|E_C\|_{L_2} \leq C \|\rho\|_{L_{6/5}} , \quad (5.12)$$

and by definition of ρ and using Holder inequality, we have

$$\begin{aligned} \|\rho\|_{L_{6/5}}^2 &\leq \|EA\|_{L_{6/5}}^2 + \|X\pi\|_{L_{6/5}}^2 + \|E\partial_s h\|_{L_{6/5}}^2 + \|\mathcal{F}\partial_s k\|_{L_{6/5}}^2 + \|kA\mathcal{F}\|_{L_{6/5}}^2 \\ &\leq \|E\|_{L_2}^2 \|A\|_{L_3}^2 + \|X\|_{L_3}^2 \|\pi\|_{L_2}^2 + \|E\|_{L_2}^2 \|\partial_s h\|_{L_3}^2 + \|\partial_s k\|_{L_3}^2 \|\mathcal{F}\|_{L_2}^2 + \|kA\|_{L_3}^2 \|\mathcal{F}\|_{L_2}^2 \\ &\leq \|E\|_{L_2}^2 \|A\|_{H_1}^2 + \|X\|_{H_1}^2 \|\pi\|_{L_3}^2 + \|E\|_{L_2}^2 \|\partial_s h\|_{H_1}^2 + \|\mathcal{F}\|_{L_2}^2 (\|\partial_s k\|_{H_1}^2 + \|kA\|_{H_1}^2) , \end{aligned} \quad (5.13)$$

where we used the Sobolev inequality to show that $\|u\|_{L_3} \leq C\|u\|_{H_1}$.

Since $\partial_s E_C^s = 4\pi\rho$, by taking Fourier transform on both sides, we have

$$\hat{E}_C^s = -i4\pi \frac{k^s}{|k|^2} \hat{\rho} , \quad (5.14)$$

hence, we have

$$\begin{aligned} \|D^2 E_C\|_{L_2}^2 &= \int_{\mathbb{R}^3} d^3x (\partial_q \partial_r E_{Cs})^2 \\ &= \int_{\mathbb{R}^3} d^3k \left| k^2 \hat{E}_{Cs} \right|^2 \\ &= (4\pi)^2 \int_{\mathbb{R}^3} d^3k |k_s \hat{\rho}|^2 \\ &= (4\pi)^2 \int_{\mathbb{R}^3} d^3x (\partial_s \rho)^2 = (4\pi)^2 \|\partial_s \rho\|_{L_2}^2 . \end{aligned} \quad (5.15)$$

Using definition of Sobolev norm and Schauder ring property, we have

$$\begin{aligned} \|\partial_s \rho\|_{L_2}^2 &\leq \|EA\|_{H_1}^2 + \|X\pi\|_{H_1}^2 + \|E\partial_s h\|_{H_1}^2 + \|\mathcal{F}\partial_s k\|_{H_1}^2 + \|kA\mathcal{F}\|_{H_1}^2 \\ &\leq \|E\|_{H_1}^2 \|A\|_{H_1}^2 + \|X\|_{H_1}^2 \|\pi\|_{H_1}^2 + \|E\|_{H_1}^2 \|\partial_s h\|_{H_1}^2 + \|\mathcal{F}\|_{H_1}^2 (\|\partial_s k\|_{H_1}^2 + \|kA\|_{H_1}^2) . \end{aligned} \quad (5.16)$$

By definition of field strength, we get an estimate

$$\|\mathcal{F}\|_{H_1} \leq C (\|A\|_{H_2} + \|A\|_{H_2}^2) , \quad (5.17)$$

and using condition (5.9),

$$\begin{aligned} \|\partial_s h\|_{H_1} &\leq \|\partial_i h\|_{H_1} \|\partial_s \phi\|_{H_1} , \\ &\leq C \|\phi\|_{H_2}^2 \end{aligned} \quad (5.18)$$

and the fact that gauge kinetic function is a holomorphic function, we have

$$\begin{aligned} \|E_C\|_{H_2} \leq & C \left(\|A\|_{H_2} \|E\|_{H_1} + \|\phi\|_{H_2} \|\pi\|_{H_1} + \|\phi\|_{H_2}^2 \|E\|_{H_1} + \|A\|_{H_2} \|\phi\|_{H_2}^2 \right. \\ & \left. + \|A\|_{H_2}^2 \|\phi\|_{H_2}^2 + \|A\|_{H_2}^3 \|\phi\|_{H_2} + \|A\|_{H_2}^2 \|\phi\|_{H_1}^2 \right), \end{aligned} \quad (5.19)$$

which shows that E_C lies in H_2 for all $u \in \mathcal{H}$ and the first part of the lemma is proven.

Now we will prove the Lipshitz condition for E_C which is important for proving the local existence of the evolution equation. From the definition of E_C we have

$$\begin{aligned} E_C^{as}(u') - E_C^{as}(u) = & \frac{\hat{r}^s}{4\pi r^2} * \left(-f_{bf}^c \{ h'^{ab} h'_{cd} A_r'^f E_d'^r - h^{ab} h_{cd} A_r^f E_d^r \} \right. \\ & - h'^{ab} g'_{ij} \{ X_b'^i \bar{\pi}'^{\bar{j}} + \bar{X}_b'^{\bar{j}} \pi'^i \} + h^{ab} g_{ij} \{ X_b^i \bar{\pi}^{\bar{j}} + \bar{X}_b^{\bar{j}} \pi^i \} \\ & + h'^{ab} (k'_{de} f_{bc}^d - k'_{bd} f_{ce}^d) A'^{cr} \tilde{\mathcal{F}}'^{es0} - h^{ab} (k_{de} f_{bc}^d - k_{bd} f_{ce}^d) A^{cr} \tilde{\mathcal{F}}^{es0} \\ & \left. - h'^{ab} E'^{cs} \partial_s h'_{bc} + h^{ab} E^{cs} \partial_s h_{bc} + h'^{ab} \tilde{\mathcal{F}}'^{cs0} \partial_s k'_{bc} - h^{ab} \tilde{\mathcal{F}}^{cs0} \partial_s k_{bc} \right). \end{aligned} \quad (5.20)$$

Recalling the estimate from the prove of the first part of lemma and using assumption that $\partial_s h_{ab}$ is locally Lipshitz, then we have

$$\begin{aligned} \|E_C^{as}(u') - E_C^{as}(u)\|_{H_2} \leq & K \left\{ \|A' - A\|_{H_2} [\|E'\|_{H_1} + \|\phi\|_{H_2}^2 (\|A\|_{H_2} + \|A\|_{H_2} \|A' + A\|_{H_2} \right. \\ & \left. + \|A\|_{H_2}^2 + \|A' + A\|_{H_2})] + \|E' - E\|_{H_1} (\|A\|_{H_2} + \|\phi\|_{H_2}^2) \right. \\ & \left. + \|\pi' - \pi\|_{H_1} \|\phi\|_{H_2} + \|\phi' - \phi\|_{H_2} [\|E\|_{H_1} \|\phi' + \phi\|_{H_2} \right. \\ & \left. + \|A\|_{H_2} (1 + \|A\|_{H_2}) \|\phi' + \phi\|_{H_2} + \|A'\|_{H_2}^2 (1 + \|A'\|_{H_2})] \right\} \\ = & C_1(\|u'\|, \|u\|) \|A' - A\|_{H_2} + C_2(\|u'\|, \|u\|) \|E' - E\|_{H_1} \\ & + C_3(\|u'\|, \|u\|) \|\phi' - \phi\|_{H_2} + C_4(\|u'\|, \|u\|) \|\pi' - \pi\|_{H_1}. \end{aligned} \quad (5.21)$$

Then for $u \in \mathcal{H}$, E_C is a locally Lipshitz function which proves the second part of the lemma. $\square$

The final proof of Lipshitz condition for J is established by showing that the other components of J are also Lipshitz functions. Using Sobolev inequality, we have

$$\begin{aligned} \|J_2(u') - J_2(u)\|_{H_1} \leq & C \left\{ \|A' - A\|_{H_2} [\|A\|_{H_2} + \|A\|_{H_2}^2 + \|E\|_{H_1} \|\phi'\|_{H_2}^2 + \|\phi\|_{H_2}^2 + \|E\|_{H_1} \|\phi'\|_{H_2}^2 \right. \\ & \left. + \|A' + A\|_{H_2} (1 + \|A'\|_{H_2} + \|\phi\|_{H_2}^2 + \|\phi'\|_{H_2} \|\pi'\|_{H_1}) + \|\phi'\|_{H_2} \|\pi'\|_{H_1} \right] \\ & + \|E' - E\|_{H_1} [\|\phi'\|_{H_2} \|\pi'\|_{H_1} + \|\phi\|_{H_2}^2 + \|A'\|_{H_2} \|\phi'\|_{H_2}^2] \\ & + \|\phi' - \phi\|_{H_2} [\|\phi'\|_{H_2} + \|\phi' + \phi\|_{H_2} (\|E\|_{H_1} + \|A'\|_{H_2} + \|A'\|_{H_2}^2 + \|A\|_{H_2}) \\ & + \|\pi\|_{H_1} \|E\|_{H_1} + \|\pi\|_{H_2} (\|A\|_{H_2} + \|A\|_{H_2}^2) + \|\phi\|_{H_2}] \\ & \left. + \|\pi' - \pi\|_{H_1} [\|E\|_{H_1} + \|\phi'\|_{H_2} (\|A\|_{H_2} + \|A\|_{H_2}^2)] \right\} \\ = & C_5(\|u'\|, \|u\|) \|A' - A\|_{H_2} + C_6(\|u'\|, \|u\|) \|E' - E\|_{H_1} \\ & + C_7(\|u'\|, \|u\|) \|\phi' - \phi\|_{H_2} + C_8(\|u'\|, \|u\|) \|\pi' - \pi\|_{H_1}, \end{aligned} \quad (5.22)$$

and

$$\begin{aligned}
\|J_4(u') - J_4(u)\|_{H_1} \leq & C \left\{ \|\Gamma' - \Gamma\|_{H_2} \left[\|\phi'\|_{H_2} + \|A'\|_{H_2} \|\phi'\|_{H_2}^2 + \|A'\|_{H_2}^2 \|\phi'\|_{H_2}^2 + \|\pi\|_{H_1}^2 \right] \right. \\
& \|E' - E\|_{H_1} \left[\|\phi\|_{H_2} \|E' + E\|_{H_1} + \|\phi'\|_{H_2} \|\mathcal{F}\|_{H_1} \right] + \|\pi' - \pi\|_{H_1} \|\Gamma\|_{H_2} \\
& \|A' - A\|_{H_2} \left[\|\phi'\|_{H_2} + \|A' - A\|_{H_2} (\|\phi'\|_{H_2} + \|\Gamma\|_{H_2} \|\phi\|_{H_2}^2) + \|\Gamma\|_{H_2} \|\phi\|_{H_2}^2 \right. \\
& \left. \left. + (1 + \|A' + A\|_{H_2}) (\|\phi\|_{H_2} \|\mathcal{F}' - \mathcal{F}\|_{H_1} + \|\phi\|_{H_2} \|E'\|_{H_1}) \right] \right. \\
& + \|\phi' - \phi\|_{H_2} \left[\|\Gamma\|_{H_2} \|\phi' + \phi\|_{H_2} (1 + \|A\|_{H_2}^2) + \|A'\|_{H_2} + \|A\|_{H_2} \right. \\
& \left. \left. + \|A\|_{H_2}^2 + \|E\|_{H_2} \|\mathcal{F}\|_{H_1} + \|E\|_{H_2}^2 + \|\mathcal{F}\|_{H_2}^2 \right] + \|\partial_j V'_S - \partial_j V_S\|_{H_1} \right\}. \tag{5.23}
\end{aligned}$$

Using estimate (4.2) and (5.17) and using the assumption in (2.5), we have

$$\begin{aligned}
\|J_4(u') - J_4(u)\|_{H_1} \leq & C_9(\|u'\|, \|u\|) \|A' - A\|_{H_2} + C_{10}(\|u'\|, \|u\|) \|E' - E\|_{H_1} \\
& + C_{11}(\|u'\|, \|u\|) \|\phi' - \phi\|_{H_2} + C_{12}(\|u'\|, \|u\|) \|\pi' - \pi\|_{H_1} \tag{5.24}
\end{aligned}$$

Thus, from lemma 5, (5.22) and (5.24), the nonlinear operator J is a mapping from $\mathcal{H}$ to itself and satisfies locally Lipschitz condition.

Then by Segal's theorem, for any initial data u_0 in $\mathcal{H}$, there exists a positive constant $T > 0$ depending on u_0 such that the equation (3.35) admits a unique mild solution which is continuous in $\mathcal{H}$ for an interval $[0, T]$, i.e. $u \in C([0, T], \mathcal{H})$ which satisfies (5.5). Furthermore, the solutions can be extended into maximal mild solutions on interval $[0, T_{\max})$ such that either

1. $T_{\max} = +\infty$ and the equation (3.35) admits a global solution, or
2. $\|u(t)\| \rightarrow \infty$ as $t \rightarrow T_{\max}$ and the solution blow up on a finite time $T_{\max}$.

If the initial value u_0 lies in $D_{\mathcal{A}}$ then equation (3.35) admits a unique classical solutions $u(t)$ for an interval $[0, T_{\max})$ which remains in $D_{\mathcal{A}}$ and satisfies differential equations

$$\frac{du(t)}{dt} = \mathcal{A}u(t) + J(u(t)), \tag{5.25}$$

with $\frac{du(t)}{dt}$ is a continuous curve in $\mathcal{H}$. Then, the solutions u belong to,

$$u \in C^1([0, T_{\max}), \mathcal{H}) \cap C([0, T_{\max}), D_{\mathcal{A}}), \tag{5.26}$$

such that either $T_{\max} = +\infty$ or $\|u(t)\| \rightarrow \infty$ as $t \rightarrow T_{\max}$.

Now, we shall show that the solution of modified equation (3.35) which satisfy the constraint is the solution of original equation.

Following result of [4] for the constraint equation, we have

$$\|\mathcal{C}(t)\|_{L_2}^2 \leq \|\mathcal{C}(0)\|_{L_2}^2 \exp \left(\int_0^t C' \|E(s)\|_{H_1} \right). \tag{5.27}$$

Since the initial value of the constraint equation is $\mathcal{C}(0) = 0$, then $\mathcal{C}(t)$ is vanished in the interval existence of $u(t)$. Hence, the solutions of modified equation (3.35) always satisfy the constraint equation then it is the solutions of the original equation.

Therefore, we have proven,

Theorem 1. *Let u_0 be any initial data lying in $\mathcal{H} = (H_2 \times H_1)^3$. If the conditions (2.5), (4.1), (4.2), and (4.3) are satisfied, then there exists a positive constant $T_{\max} > 0$ depending on u_0 such that the integral equation (5.5) admits a unique maximal solution $u(t)$ on interval $[0, T_{\max})$ which belongs to $u \in C([0, T_{\max}), \mathcal{H})$ and either*

1. $T_{\max} = +\infty$ and the equation (3.35) admits a global solution, or
2. $\|u(t)\| \rightarrow \infty$ as $t \rightarrow T_{\max}$ and the solution blow up on a finite time $T_{\max}$.

Furthermore, if u_0 lies in $D_{\mathcal{A}}$ and satisfies the constraint $\mathcal{C}(u_0) = 0$, then the equation (3.35) admits a unique classical solution $u(t)$ for an interval $[0, T_{\max})$ which remains in $D_{\mathcal{A}}$, and belong to $u \in C^1([0, T_{\max}), \mathcal{H}) \cap C([0, T_{\max}), D_{\mathcal{A}})$, and satisfy the constraint $\mathcal{C}(u(t)) = 0$ such that either $T_{\max} = +\infty$ or $\|u(t)\| \rightarrow \infty$ as $t \rightarrow T_{\max}$.

A Convention and Notation

The purpose of this appendix is to inform our conventions used in this paper. The spacetime metric is flat with the signature $(-, +, +, +)$.

The following indices are used:

$\mu, \nu, \rho, \sigma = 0, \dots, 3,$	label 4-dimensional flat spacetime
$r, s, p, q = 1, 2, 3,$	label 3-dimensional flat space
$i, \bar{i}, j, \bar{j}, k, \bar{k} = 1, \dots, n_c,$	label n_c dimensional Kähler manifold
$a, b, c, d = 1, \dots, n_v,$	label the gauge index

B List of Inequality

In this appendix, we mention some basic inequalities used in this paper. For detail reviews, see [10, 12].

Young inequality for convolution

Suppose $f \in L_p(\mathbb{R}^d)$ and $g \in L_q(\mathbb{R}^d)$ and

$$\frac{1}{p} + \frac{1}{q} = \frac{1}{r} + 1, \quad (\text{B.1})$$

with $p, q, r \in \mathbb{R}$ and $1 \leq p, q, r \leq \infty$. Then

$$\|f * g\|_{L_r} \leq \|f\|_{L_p} \|g\|_{L_q}. \quad (\text{B.2})$$

If $g \in L_{q,w}(\mathbb{R}^d)$ where $L_{q,w}$ is weak L_q space, then

$$\|f * g\|_{L_r} \leq \|f\|_{L_p} \|g\|_{L_{q,w}}, \quad (\text{B.3})$$

with p, q, r are defined and satisfy condition above.

Holder inequality

Let p, q, r be positive numbers and satisfy $p, q, r \leq 1$ and

$$\frac{1}{p} + \frac{1}{q} = \frac{1}{r}. \quad (\text{B.4})$$

Suppose $f \in L_p(\mathbb{R}^d)$ and $g \in L_q(\mathbb{R}^d)$, then $fg \in L_r(\mathbb{R}^d)$ and

$$\|fg\|_{L_r} \leq \|f\|_{L_p} \|g\|_{L_q}. \quad (\text{B.5})$$

Sobolev inequality

Let q, m be real numbers with $1 \leq q, m \leq \infty$ and r, s are integers where $0 \leq r < s$ which satisfy

$$\frac{1}{p} = \frac{r}{3} + \theta \left(\frac{1}{m} - \frac{s}{3} \right) + (1 - \theta) \frac{1}{q}, \quad (\text{B.6})$$

with $r/s \leq \theta \leq 1$ and p is non negative. For $u \in H_s(\mathbb{R}^3) \cap L_q(\mathbb{R}^3)$, there is a positive constant C which depends only on m, j, q, r and θ such that the following inequality holds

$$\|\partial^r u\|_{L_p} \leq C \|\partial^s u\|_{L_m}^\theta \|u\|_{L_q}^{1-\theta}. \quad (\text{B.7})$$

Acknowledgments

Our research is supported by Hibah Kompetensi DIKTI 2012 No. 781a/I1.C01/PL/2012 and Riset Desentralisasi DIKTI-ITB 2012 No.003.8/TL-J/DIPA/SPK/2012.

References

- [1] Segal, I. (1963), "Nonlinear Semigroup", *Ann. Math.*, 78, 339.
- [2] Segal, I. (1979), "The Cauchy Problem for the Yang-Mills Equations", *J. Func. Anal.*, 33, 175.
- [3] Ginibre, J. and Velo, G. (1981), "The Cauchy Problem for Coupled Yang-Mills and Scalar Fields in the Temporal Gauge", *Commun. Math. Phys.*, 82, 1-28.
- [4] Eardley, D. and Moncrief, V. (1982), "The Global Existence of Yang-Mills-Higgs Fields in 4-Dimensional Minkowski Space : I. Local Existence and Smoothness Properties", *Commun. Math. Phys.*, 83, 171-191.
- [5] Eardley, D. and Moncrief, V. (1982), "The Global Existence of Yang-Mills-Higgs Fields in 4-Dimensional Minkowski Space : II. Completion of Proof", *Commun. Math. Phys.*, 83, 193-212.
- [6] Wess, J and Bagger, J. (1992), *Supersymmetry and Supergravity 2nd edition*, Princeton University Press
- [7] D'Auria, R. and Ferrara, S. (2001), "On Fermion Masses, Gradient Flows and Potential in Supersymmetric Theories", *JHEP* **0105**, 034, hep-th/0103153.

- [8] Andrianopoli, L., D'Auria, R., and Ferrara, S. (2002), "Supersymmetry reduction of N -extended supergravities in four dimensions", *JHEP* **0203**, 025; hep-th/0110277.
- [9] Akbar, F.T., Gunara, B., Zen, F. and Triyanta (2012), "Local Existence of $N=1$ Supersymmetric Gauge Theory in Four Dimensions", *AIP Conference Proceeding of Asian Physics Symposium*.
- [10] Reed, M and Simmon, B. (1980), *Methods of Modern Mathematical Physics vol I and II*, Academic Press.
- [11] S. Zheng (2004), *Nonlinear Evolution Equations*, Chapman and Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, 133.
- [12] Adams, Robert (1975), *Sobolev Spaces*, Boston, MA: Academic Press.